

dd-sequences and partial Euler-Poincaré characteristics of Koszul complex

Nguyen Tu Cuong* and Doan Trung Cuong†

Institute of Mathematics

18 Hoang Quoc Viet Road, 10307 Hanoi, Vietnam

March 23, 2005

Abstract

The aim of this paper is to introduce a new notion of sequences called dd-sequences and show that this notion may be convenient for studying the polynomial property of partial Euler-Poincaré characteristics of the Koszul complex with respect to the powers of a system of parameters. Some results about the dd-sequences, the partial Euler-Poincaré characteristics and the lengths of local cohomology modules are also presented in the paper. There are also applications of dd-sequences on the structure of sequentially Cohen-Macaulay modules.

Keywords: dd-sequence, p-standard system of parameters, Koszul complex, Euler-Poincaré characteristic, local cohomology, sequentially Cohen-Macaulay.

AMS Classification: 13D99, 13D45, 13H15

1 Introduction

Let $(R, \mathfrak{m})$ be a commutative local Noetherian ring and M a finitely generated R -module of dimension d . Let $\underline{x} = (x_1, \dots, x_d)$ be a system of parameters of M . Denote by $H_i(\underline{x}; M)$ the i -th Koszul homology module of M with respect to the system of parameters $\underline{x}$. Following Serre [21, Appendice II], the k -th Euler-Poincaré characteristic of M with respect to $\underline{x}$ is defined by

$$\chi_k(\underline{x}; M) = \sum_{i=k}^{\infty} (-1)^{i-k} \ell(H_i(\underline{x}; M))$$

for $k = 0, 1, \dots, d$, where $\ell(N)$ stands for the length of the R -module N . Let $\underline{n} = (n_1, \dots, n_d)$ be a d -tuple of positive integers. We denote by $\underline{x}(\underline{n})$ the system of parameters $(x_1^{n_1}, \dots, x_d^{n_d})$. Then we can consider $\chi_k(\underline{x}(\underline{n}); M)$ as a function

*Email: ntcuong@math.ac.vn

†Email: dtcuong@math.ac.vn

in $\underline{n}$. It is well-known by Garcia Roig [11] that this function in general is not a polynomial. But V. T. Khoi and the first author have proved in [8] that the least degree of all polynomials in $\underline{n}$ bounding above the function $\chi_k(\underline{x}(\underline{n}); M)$ is independent of the choice of the system of parameters $\underline{x}$ and they denote this invariant by $p_k(M)$. It was shown in [7] (in that paper $\chi_1(\underline{x}(\underline{n}); M)$ and $p_1(M)$ are denoted by $I_M(\underline{n}; \underline{x})$ and $p(M)$ respectively) that the 1-st Euler-Poincaré characteristic of M is a polynomial in $\underline{n}$ provided $\underline{x}$ is a p-standard system of parameters of M . Moreover, in this case the polynomial has a very simple formula as follows

$$\chi_1(\underline{x}(\underline{n}); M) = \sum_{i=0}^{p_1(M)} n_1 \dots n_i e(x_1, \dots, x_i; (0 : x_{i+1})_{M/(x_{i+2}, \dots, x_d)M}),$$

where, for a system of parameters $\underline{y}$ of a R -module N , we denote by $e(\underline{y}; N)$ the Serre multiplicity of N with respect to $\underline{y}$ and set

$$e(x_1, \dots, x_i; (0 : x_{i+1})_{M/(x_{i+2}, \dots, x_d)M}) = \ell((x_2, \dots, x_d)M : x_1/(x_2, \dots, x_d)M)$$

for $i = 0$. Recall that a system of parameters $\underline{x} = (x_1, \dots, x_d)$ of M is called a p-standard system of parameters if $x_i \in \mathfrak{a}(M/(x_{i+1}, \dots, x_d)M)$ for $i = 1, \dots, d$, where $\mathfrak{a}(M) = \mathfrak{a}_0(M) \dots \mathfrak{a}_{d-1}(M)$ and $\mathfrak{a}_i(M)$ is the annihilator ideal of the i -th local cohomology module $H_{\mathfrak{m}}^i(M)$ with respect to the maximal ideal $\mathfrak{m}$. It should be mentioned that the notion of p-standard system of parameters has been used as an important tool for the resolution of Macaulayfication by T. Kawasaki [16, 17]. Therefore it raises to a natural question: Is $\chi_k(\underline{x}(\underline{n}); M)$ still a polynomial in $\underline{n}$ for all $k > 1$ provided $\underline{x}$ is a p-standard system of parameters?

In order to give an affirmative answer to the above question we introduce in this paper the concept of dd-sequence, which is a slight generalization of the notion of p-standard sequence. Note here that while there does not exist a necessary and sufficient condition for a system of parameters to be p-standard, we can characterize a system of parameters to be a dd-sequence in term of the 1-st Euler-Poincaré characteristic. Therefore we can solve the above question for the class of systems of parameters which are dd-sequences. Moreover, as an application, we can study more details about systems of parameters, which are dd-sequences, as well as the partial Euler-Poincaré characteristics with respect to them.

Let us to summarize the main results of this paper. The paper is divided into 6 sections. We need first a necessary and sufficient condition for the k -th Euler-Poincaré characteristic $\chi_k(\underline{x}(\underline{n}); M)$ of M to be a polynomial in $\underline{n}$. Hence the following theorem, which is a generalization of a result of [4], is the main result of Section 2.

Theorem 1.1. *Let M be a finitely generated R -module, $\underline{x} = (x_1, \dots, x_d)$ a system of parameters of M and n_0 a positive integer. Then the k -th Euler-Poincaré characteristic $\chi_k(\underline{x}(\underline{n}); M)$ is a polynomial in $\underline{n} = (n_1, \dots, n_d)$ for all $n_i \geq n_0$ if and only if the following condition is satisfied*

$$(0 : x_i^{n_i})_{H_{k-1}(x_1^{n_1}, \dots, x_{i-1}^{n_{i-1}}, x_{i+1}^{n_{i+1}}, \dots, x_d^{n_d}; M)} = (0 : x_i^{n_0})_{H_{k-1}(x_1^{n_1}, \dots, x_{i-1}^{n_{i-1}}, x_{i+1}^{n_{i+1}}, \dots, x_d^{n_d}; M)}$$

for all $i = 1, \dots, d$ and $n_1, \dots, n_d \geq n_0$.

Next, in Section 3 we use the concept of d-sequence of Huneke (see [15]) to introduce the notion of dd-sequences. A sequence $\underline{x} = (x_1, \dots, x_s)$ of elements in $\mathfrak{m}$ is said to be a dd-sequence of M if for all $i \in \{1, \dots, s\}$ and all s -tuples of positive integers $(n_1, \dots, n_s)$, the sequence $x_1^{n_1}, \dots, x_i^{n_i}$ is a d-sequence of the module $M/(x_{i+1}^{n_{i+1}}, \dots, x_s^{n_s})M$. Then the main result of this section is the following characterization of a system of parameters to be a dd-sequence.

Theorem 1.2. *Let M be a finitely generated R -module and $\underline{x} = (x_1, \dots, x_d)$ a system of parameters of M . Then $\underline{x}$ is a dd-sequence if and only if*

$$\chi_1(\underline{x}(\underline{n}); M) = \sum_{i=0}^{p_1(M)} n_1 \dots n_i e(x_1, \dots, x_i; (0 : x_{i+1})_{M/(x_{i+2}, \dots, x_d)M})$$

for all $n_1, \dots, n_d > 0$.

It follows from this theorem that every p-standard system of parameters of M is a dd-sequence. However, the converse statement is wrong. Observe that there always exist p-standard systems of parameters of M provided the ring R admits a dualizing complex (see [6]). Hence, in this case, there exist also systems of parameters of M which are dd-sequences.

The Section 4 is devoted to show the following theorem, which is one of the main results of the paper.

Theorem 1.3. *Let M be a finitely generated R -module, $\underline{x} = (x_1, \dots, x_d)$ a system of parameters of M . Suppose that $\underline{x}$ is a dd-sequence of M . Then the k -th Euler-Poincaré characteristic $\chi_k(\underline{x}(\underline{n}); M)$ is a polynomial in $\underline{n}$ for all $k > 0$. Moreover, in this case we have*

$$\chi_k(\underline{x}(\underline{n}); M) = \sum_{i=0}^{p_k(M)} n_1 \dots n_i e(x_1, \dots, x_i; (0 : x_{i+1})_{H_{k-1}(x_{i+2}, \dots, x_d; M)}),$$

where we set $e(x_1, \dots, x_i; (0 : x_{i+1})_{H_{k-1}(x_{i+2}, \dots, x_d; M)}) = \ell((0 : x_1)_{H_{k-1}(x_2, \dots, x_d; M)})$ for $i = 0$.

Let k be an integer with $k \geq p_1(M)$ and $\underline{x} = (x_1, \dots, x_d)$ a system of parameters which is a dd-sequence. Then it follows from Theorem 1.2 that $p_1(M/(x_1, \dots, x_k)M) \leq 0$. Therefore $M/(x_1, \dots, x_k)M$ is a generalized Cohen-Macaulay module of dimension $d - k$. Thus $\ell(H_{\mathfrak{m}}^i(M/(x_1, \dots, x_k)M)) < \infty$ for all $i = 0, \dots, d - k - 1$.

As an application of Theorem 1.3, we prove in Section 5 a result concerning with the length of local cohomology modules as follows.

Theorem 1.4. *Let M be a finitely generated R -module and $\underline{x}$ a system of parameters of M . Assume that $\underline{x}$ is a dd-sequence. Then $\ell(H_{\mathfrak{m}}^i(M/(x_1^{n_1}, \dots, x_k^{n_k})M))$ is a polynomial in $\underline{n}$ for all $k \geq p_1(M)$ and $i < d - k$.*

The last Section is to focus on sequentially Cohen-Macaulay modules which were introduced first for graded algebras by Stanley [22]. We show that there always exists system of parameters which is a dd-sequence on a sequentially Cohen-Macaulay module and their partial Euler-Poincaré characteristic are easy to compute by multiplicities of the dimension filtration of the module. Therefore the following result is the main result of this section.

Theorem 1.5. *Let $\underline{x} = (x_1, \dots, x_d)$ be a system of parameters of a sequentially Cohen-Macaulay module M . Suppose that*

$$M_0 \subset M_1 \subset \dots \subset M_{t-1} \subset M_t = M$$

is the dimension filtration of M with $\dim M_i = d_i$. Then it holds

(a) The following statements are equivalent:

- i) $M_i \cap (x_{d_i+1}, \dots, x_d)M = 0$ for all $i = 0, 1, \dots, t-1$.*
- ii) $(x_{d_i+1}, \dots, x_d)M_i = 0$ for all $i = 0, 1, \dots, t-1$.*
- iii) For all $n_1, \dots, n_d \geq 1$,*

$$\chi_1(\underline{x}(\underline{n}); M) = \sum_{i=0}^{t-1} n_1 \dots n_{d_i} e(x_1, \dots, x_{d_i}; M_i).$$

iv) $\underline{x}$ is a dd-sequence of M .

(b) Suppose that $\underline{x}$ is satisfied one of the equivalent conditions above, then

$$\chi_k(\underline{x}(\underline{n}); M) = \sum_{d_i \leq d-k} a_i n_1 \dots n_{d_i} e(x_1, \dots, x_{d_i}; M_i),$$

where

$$a_{t-i} = \sum_{\substack{j_1 + \dots + j_{i-1} \leq k-1 \\ k \leq j_1 + \dots + j_i}} (-1)^{j_1 + \dots + j_i - k} \binom{d_t - d_{t-1}}{j_1} \dots \binom{d_{i+1} - d_i}{j_{t-i}},$$

and we stipulate that $\binom{a}{b} = 0$ if $a < b$.

2 Partial Euler-Poincaré characteristics

Throughout this paper we always denote by $(R, \mathfrak{m})$ a commutative Noetherian local ring with the maximal ideal $\mathfrak{m}$ and by M a finitely generated R -module of dimension d . Let $\underline{x} = (x_1, \dots, x_d)$ be a system of parameters of M . Consider the Koszul complex $K(\underline{x}; M)$ of M with respect to the system of parameters $\underline{x}$. Let $H_i(\underline{x}; M)$ be the i -th Koszul homology module of $K(\underline{x}; M)$. Then the k -th Euler-Poincaré characteristic of M with respect to $\underline{x}$ (or the partial Euler-Poincaré characteristic of $K(\underline{x}; M)$) is defined by

$$\chi_k(\underline{x}; M) = \sum_{i=k}^{\infty} (-1)^{i-k} \ell(H_i(\underline{x}; M)).$$

It is well-known that $\chi_0(\underline{x}; M) = e(\underline{x}; M)$, $\chi_1(\underline{x}; M) = \ell(M/\underline{x}M) - e(\underline{x}; M)$, $\chi_k(\underline{x}; M) \geq 0$ for all k and $\chi_k(\underline{x}; M) = 0$ for all $k > d$. The following lemma is a generalization of a result by M. Auslander and D. A. Buchsbaum (see [1, Corollary 4.3]) and is often used in the sequel.

Lemma 2.1. [8, Corollary 2.2] *Let $\underline{x} = (x_1, \dots, x_d)$ be a system of parameters of M . Then*

$$\chi_k(\underline{x}; M) = \sum_{i=0}^{d-k} e(x_1, \dots, x_i; (0 : x_{i+1})_{H_{k-1}(x_{i+2}, \dots, x_d; M)}).$$

Let now $\underline{n} = (n_1, \dots, n_d)$ be a d-tuple of positive integers. Put $\underline{x}(\underline{n}) = (x_1^{n_1}, \dots, x_d^{n_d})$ and consider $\chi_k(\underline{x}(\underline{n}); M)$ as a function in $\underline{n}$.

In order to prove Theorem 1.1 we need the following auxiliary lemmata.

Lemma 2.2. *Let $\underline{x} = (x_1, \dots, x_d)$ be a system of parameters of M . Then the following statements are true for all $k \geq 0$.*

- (i) $\chi_k(\underline{x}(\underline{n}); M) \leq n_1 \dots n_d \chi_k(\underline{x}; M)$ for all $\underline{n}$.
- (ii) $\chi_k(\underline{x}(\underline{n}); M)$ is an increasing function, i. e., for all $\underline{m} = (m_1, \dots, m_d)$, $n_1 \leq m_1, \dots, n_d \leq m_d$,

$$\chi_k(\underline{x}(\underline{n}); M) \leq \chi_k(\underline{x}(\underline{m}); M).$$

- (iii) *If $\chi_k(\underline{x}(\underline{n}); M)$ is a polynomial in $\underline{n}$ then it is linear in each variable n_i , i. e., there exist integers $\lambda_{i_1 \dots i_t}$ such that*

$$\chi_k(\underline{x}(\underline{n}); M) = \sum_{t=0}^d \sum_{0 < i_1 < \dots < i_t \leq d} \lambda_{i_1 \dots i_t} n_{i_1} \dots n_{i_t},$$

where we set $\lambda_{i_1 \dots i_t} n_{i_1} \dots n_{i_t} = \lambda$ the constant coefficient of the polynomial when $t = 0$.

Proof. The statements (i), (ii) were proved in [8, Lemma 3.2] and (iii) follows easily from (i). $\square$

Remark 2.3. Garcia Roig and Kirby proved in [12] that the function $\chi_k(\underline{x}(\underline{n}); M)$ in general is not a polynomial. But, it was shown in [11] that for the case $n_1 = n_2 = \dots = n_d = n$ the function $\chi_k(\underline{x}(n); M)$, considered as a function in one variable n , is always bounded above by a polynomial in n of degree at most $d - k$. Later, a more general result has been proved in [8], which says that the least degree of all polynomials in $\underline{n}$ bounding above the function $\chi_k(\underline{x}(\underline{n}); M)$ is independent of the choice of the system of parameters $\underline{x}$. Denote this new invariant by $p_k(M)$. Then we have

$$p_0(M) = d > p_1(M) \geq \dots \geq p_{d-1}(M) \geq p_d(M).$$

Especially, the invariant $p_1(M)$ was called in [6] the polynomial type of M and denoted by $p(M)$.

The next lemma easily follows from properties of linear polynomials.

Lemma 2.4. *Let $\varphi(n_1, \dots, n_d) : \mathbb{N}^d \longrightarrow \mathbb{N} \cup \{0\}$ be a function defining on $\mathbb{N}^d$ ($\mathbb{N}$ is the set of all positive integers). Assume that $\varphi(n_1, \dots, n_d)$ is a linear polynomial in each variable n_i for all $i = 1, \dots, d$. Then $\varphi(n_1, \dots, n_d)$ is a polynomial in $\underline{n}$.*

Proof of Theorem 1.1.

Necessary condition: Assume that $\chi_k(\underline{x}(\underline{n}); M)$ is a polynomial for all positive integers $n_1, \dots, n_d \geq n_0$. Since the function $\chi_k(\underline{x}(\underline{n}); M)$ does not depend on the order of the sequence $\underline{x}$, we need only to check the necessary condition for the case $i = d$. Using Lemma 2.1 to the sequence $x_d, x_{d-1}, \dots, x_1$ we have

$$\begin{aligned} \chi_k(\underline{x}(\underline{n}); M) &= n_d \sum_{i=k}^{d-1} n_{i+1} \dots n_{d-1} e(x_{i+1}, \dots, x_d; (0 : x_i^{n_i})_{H_{k-1}(x_1^{n_1}, \dots, x_{i-1}^{n_{i-1}}; M)}) \\ &\quad + \ell((0 : x_d^{n_d})_{H_{k-1}(x_1^{n_1}, \dots, x_{d-1}^{n_{d-1}}; M)}). \end{aligned}$$

Since $\chi_k(\underline{x}(\underline{n}); M)$ is a polynomial for all $n_i \geq n_0$ by the hypothesis, $\chi_k(\underline{x}(\underline{n}); M)$ is linear in n_d by Lemma 2.2, (iii). Therefore $\ell((0 : x_d^{n_d})_{H_{k-1}(x_1^{n_1}, \dots, x_{d-1}^{n_{d-1}}; M)})$ is a polynomial in n_d , for all $n_i \geq n_0$. Thus

$$(0 : x_d^{n_d})_{H_{k-1}(x_1^{n_1}, \dots, x_{d-1}^{n_{d-1}}; M)} = (0 : x_d^{n_0})_{H_{k-1}(x_1^{n_1}, \dots, x_{d-1}^{n_{d-1}}; M)},$$

for all $n_1, \dots, n_d \geq n_0$.

Sufficient condition: Without loss of generality we may assume that $n_0 = 1$. For each $i \in \{1, \dots, d\}$, applying Lemma 2.1 to the sequence $x_i^{n_i}, x_1^{n_1}, \dots, \widehat{x_i^{n_i}}, \dots, x_d^{n_d}$, where $\widehat{x}$ indicates that x is omitted there, we obtain by the hypothesis that

$$\begin{aligned} \chi_k(\underline{x}(\underline{n}); M) &= n_i \phi_i(n_1, \dots, \widehat{n_i}, \dots, n_d) + \ell((0 : x_i^{n_i})_{H_{k-1}(x_1^{n_1}, \dots, \widehat{x_i^{n_i}}, \dots, x_d^{n_d}; M)}) \\ &= n_i \phi_i(n_1, \dots, \widehat{n_i}, \dots, n_d) + \ell((0 : x_i)_{H_{k-1}(x_1^{n_1}, \dots, \widehat{x_i^{n_i}}, \dots, x_d^{n_d}; M)}), \end{aligned}$$

where the function $\phi_i(n_1, \dots, \widehat{n_i}, \dots, n_d)$ is independent of n_i . Therefore, for all given $(d-1)$ -tuple of positive integers $(n_1, \dots, \widehat{n_i}, \dots, n_d) \in \mathbb{N}^{d-1}$, the function $\chi_k(\underline{x}(\underline{n}); M)$ is a linear polynomial in n_i for all $i = 1, \dots, d$. Hence, $\chi_k(\underline{x}(\underline{n}); M)$ is a polynomial in $\underline{n}$ by Lemma 2.4. $\square$

3 dd-Sequences

First, we recall the notion of d-sequence.

Definition 3.1. Let $\underline{x} = (x_1, \dots, x_s)$ be a sequence of elements in the maximal ideal $\mathfrak{m}$. Then $\underline{x}$ is called a *d-sequence* of the R -module M if

$$(x_1, \dots, x_{i-1})M : x_j = (x_1, \dots, x_{i-1})M : x_i x_j$$

for all $i = 1, \dots, s$ and all $j \geq i$. Moreover, $\underline{x}$ is called a strong d-sequence of M if $\underline{x}(\underline{n}) = (x_1^{n_1}, \dots, x_s^{n_s})$ is a d-sequence for all $\underline{n} = (n_1, \dots, n_s) \in \mathbb{N}^s$.

It should be mentioned that the notion of d-sequence was introduced by C. Huneke [15] and it has become a useful tool in different topics of commutative algebra. Now we define a new notion of a sequence.

Definition 3.2. A sequence $\underline{x} = (x_1, \dots, x_s)$ of elements in the maximal ideal $\mathfrak{m}$ is called a *dd-sequence* of M if $\underline{x}$ is a strong d-sequence and the following inductive conditions are satisfied:

- (i) either $s = 1$, or
- (ii) $s > 1$ and the sequence $x_1, \dots, x_{s-1}$ is a dd-sequence of $M/x_s^n M$ for all positive integers n .

Remark 3.3. (i) It is easy to see that an M -sequence is always a dd-sequence. An unconditioned strong d-sequence is also a dd-sequence. Therefore a part of system of parameters of a Buchsbaum module is a dd-sequence.

(ii) If $x_1, \dots, x_s$ is a dd-sequence of M then $x_1^{n_1}, \dots, x_s^{n_s}$ is a dd-sequence of M for all s -tuples of positive integers $(n_1, \dots, n_s) \in \mathbb{N}^s$.

(iii) It is not difficult to show by the definition that a sequence $\underline{x} = (x_1, \dots, x_s)$ is a dd-sequence of M if and only if $x_1^{n_1}, \dots, x_i^{n_i}$ is a d-sequence of the module $M/(x_{i+1}^{n_{i+1}}, \dots, x_s^{n_s})M$ for all $i \in \{1, \dots, s\}$ and all s -tuples of positive integers $(n_1, \dots, n_s) \in \mathbb{N}^s$, where we set $M/(x_{i+1}^{n_{i+1}}, \dots, x_s^{n_s})M = M$ when $i = s$.

Let $x_1, \dots, x_s$ be a sequence of elements of $\mathfrak{m}$. For convenience, from now on we denote the sequence $x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_s$ by $x_1, \dots, \widehat{x}_i, \dots, x_s$.

Proposition 3.4. Let $x_1, \dots, x_s$ be a dd-sequence of M . Then $x_1, \dots, \widehat{x}_i, \dots, x_s$ is a dd-sequence of $M/x_i M$ for $i \in \{1, \dots, s\}$.

Proof. The proposition is proved by induction on s . The assertion follows immediately from the definition for cases $s = 1, 2$. Now assume that $s > 2$. By virtue of Remark 3.3 we have only to prove the assertion for $1 < i < s$. Since $x_1, \dots, x_{s-1}$ is a dd-sequence of $M/x_s M$ by the definition, it follows from the inductive hypothesis that $x_1, \dots, \widehat{x}_i, \dots, x_{s-1}$ is a dd-sequence of $M/(x_i, x_s)M$. Therefore, in order to prove that $x_1, \dots, \widehat{x}_i, \dots, x_s$ is a dd-sequence of $M/x_i M$, we have to verify that $x_1, \dots, \widehat{x}_i, \dots, x_s$ is a strong d-sequence of $M/x_i M$. Moreover, since $x_1^{n_1}, \dots, x_s^{n_s}$ is also a dd-sequence of M for all $(n_1, \dots, n_s) \in \mathbb{N}^s$ by Remark 3.3, (ii), we need only to show that $x_1, \dots, \widehat{x}_i, \dots, x_s$ is a d-sequence of $M/x_i M$. In fact, since $x_1, \dots, x_{s-1}$ is a dd-sequence of M/x_s^n for all $n > 0$, we can prove by using of Remark 3.3, (iii) and Krull's Intersection Theorem that $x_1, \dots, x_{s-1}$ is a dd-sequence of M . Therefore $x_1, \dots, \widehat{x}_i, \dots, x_{s-1}$ is a d-sequence of $M/x_i M$ by inductive hypothesis. On the other hand, since $x_2, \dots, x_s$ is a dd-sequence of $M/x_1 M$, $x_2, \dots, \widehat{x}_i, \dots, x_s$ is a dd-sequence of $M/(x_1, x_i)M$ by the inductive hypothesis. Combine these facts and keep in mind of the definition of a d-sequence, it remains to check that $0 :_{M/x_i M} x_s = 0 :_{M/x_i M} x_1 x_s$ or equivalent, $x_i M : x_s = x_i M : x_1 x_s$. First, we show that $x_i M : x_1 \subseteq x_i M : x_s$. To do this let a be an arbitrary element of $x_i M : x_1$. Then, there exists $b \in M$ such that $x_1 a = x_i b$, hence $b \in x_1 M : x_i \subseteq x_1 M : x_s$. Thus there exists an element $a' \in M$ such that $x_s b = x_1 a'$. Therefore, $x_1 x_s a = x_i x_s b = x_1 x_i a'$. It follows that

$x_s a - x_i a' \in 0 :_M x_1 \subseteq 0 :_M x_s$, and hence $x_s(x_s a - x_i a') = 0$. This leads to $x_s^2 a = x_s x_i a' \in x_i M$. So we get $a \in x_i M : x_s^2$. Since $\underline{x}$ is a dd-sequence of M , $x_i M : x_s^2 = x_i M : x_s$. Therefore $x_i M : x_1 \subseteq x_i M : x_s$. Finally, we obtain

$$\begin{aligned} x_i M : x_1 x_s &= (x_i M : x_1) : x_s \\ &\subseteq (x_i M : x_s) : x_s = x_i M : x_s^2 = x_i M : x_s \end{aligned}$$

and the proof is complete. $\square$

Corollary 3.5. *Let $x_1, \dots, x_s$ be a dd-sequence of M . Then for all $i \in \{1, \dots, s\}$, the sequence $x_1, \dots, \widehat{x_i}, \dots, x_s$ is also a dd-sequence of M .*

Proof. Since $x_1, \dots, x_s$ is a dd-sequence, $x_1, \dots, x_i^n, \dots, x_s$ is a dd-sequence of M for all $n > 0$. Hence, by Proposition 3.4, $x_1, \dots, \widehat{x_i}, \dots, x_s$ is also a dd-sequence of $M/x_i^n M$. Therefore the corollary follows easily from Remark 3.3, (iii) and Krull's Intersection Theorem. $\square$

Now we prove the main result of this Section.

Proof of Theorem 1.2.

Necessary condition: Recall that if $\chi_1(\underline{x}(\underline{n}); M)$ is a polynomial in $\underline{n}$, then this polynomial must have the degree $p_1(M) \leq d - 1$. Therefore the statement is proved if we can show that

$$\chi_1(\underline{x}(\underline{n}); M) = \sum_{i=0}^{d-1} n_1 \dots n_i e(x_1, \dots, x_i; (0 : x_{i+1})_{M/(x_{i+2}, \dots, x_d)M})$$

for all $n_1, \dots, n_d > 0$. We proceed by induction on d . If $d = 1$ there is nothing to do. Assume that $d > 1$. By Lemma 2.1 we have

$$\chi_1(\underline{x}(\underline{n}); M) = \sum_{i=1}^d n_{i+1} \dots n_d e(x_{i+1}, \dots, x_d; (0 : x_i^{n_i})_{M/(x_1^{n_1}, \dots, x_{i-1}^{n_{i-1}})M}).$$

Since $\underline{x}$ is a strong d-sequence of M ,

$$\begin{aligned} (0 : x_i^{n_i})_{M/(x_1^{n_1}, \dots, x_{i-1}^{n_{i-1}})M} &\subseteq (0 : x_d^{n_d})_{M/(x_1^{n_1}, \dots, x_{i-1}^{n_{i-1}})M} \\ &= (0 : x_d)_{M/(x_1^{n_1}, \dots, x_{i-1}^{n_{i-1}})M} \end{aligned}$$

for all $i \leq d$. Thus

$$e(x_{i+1}, \dots, x_d; (0 : x_i^{n_i})_{M/(x_1^{n_1}, \dots, x_{i-1}^{n_{i-1}})M}) = 0, \quad i < d,$$

and

$$\ell((0 : x_d^{n_d})_{M/(x_1^{n_1}, \dots, x_{d-1}^{n_{d-1}})M}) = \ell((0 : x_d)_{M/(x_1^{n_1}, \dots, x_{d-1}^{n_{d-1}})M}).$$

Therefore the function $\chi_1(\underline{x}(\underline{n}); M)$ is independent of n_d . Put $M' = M/x_d M$ and $\underline{x}'(\underline{n}') = (x_1^{n_1}, \dots, x_{d-1}^{n_{d-1}})$. Then we have

$$\begin{aligned} \chi_1(\underline{x}(\underline{n}); M) &= \chi_1(\underline{x}'(\underline{n}'), x_d; M) \\ &= \chi_1(\underline{x}'(\underline{n}'); M') + n_1 \dots n_{d-1} e(x_1, \dots, x_{d-1}; 0 :_M x_d). \end{aligned}$$

It follows by the inductive hypothesis that

$$\begin{aligned}\chi_1(\underline{x}'(\underline{n}'); M') &= \sum_{i=0}^{d-2} n_1 \dots n_i e(x_1, \dots, x_i; (0 : x_{i+1})_{M'/(x_{i+2}, \dots, x_{d-1})M'}) \\ &= \sum_{i=0}^{d-2} n_1 \dots n_i e(x_1, \dots, x_i; (0 : x_{i+1})_{M/(x_{i+2}, \dots, x_{d-1}, x_d)M}).\end{aligned}$$

Therefore

$$\chi_1(\underline{x}(\underline{n}); M) = \sum_{i=0}^{d-1} n_1 \dots n_i e(x_1, \dots, x_i; (0 : x_{i+1})_{M/(x_{i+2}, \dots, x_d)M})$$

as required.

Sufficient condition: We also prove the statement by induction on d . The case $d = 1$ is trivial. Suppose that $d > 1$. Set $\overline{M} = M/x_d^{n_d}M$ for $n_d \in \mathbb{N}$. Then we have

$$\begin{aligned}\chi_1(\underline{x}'(\underline{n}'); \overline{M}) &= \chi_1(\underline{x}(\underline{n}); M) - n_1 \dots n_{d-1} e(x_1, \dots, x_{d-1}; 0 :_M x_d^{n_d}) \\ &= \sum_{i=0}^{d-2} n_1 \dots n_i e(x_1, \dots, x_i; (0 : x_{i+1})_{M/(x_{i+2}, \dots, x_{d-1}, x_d)M}) \\ &\quad + n_1 \dots n_{d-1} (e(x_1, \dots, x_{d-1}; 0 :_M x_d) - e(x_1, \dots, x_{d-1}; 0 :_M x_d^{n_d})).\end{aligned}$$

Since $\chi_1(\underline{x}'(\underline{n}'); \overline{M})$ forms a polynomial in $\underline{n}$,

$$\deg(\chi_1(\underline{x}'(\underline{n}'); \overline{M})) = p_1(\overline{M}) \leq \dim \overline{M} - 1 = d - 2.$$

Therefore

$$e(x_1, \dots, x_{d-1}; 0 :_M x_d) - e(x_1, \dots, x_{d-1}; 0 :_M x_d^{n_d}) = 0.$$

Hence

$$\chi_1(\underline{x}'(\underline{n}'); \overline{M}) = \sum_{i=0}^{d-2} n_1 \dots n_i e(x_1, \dots, x_i; (0 : x_{i+1})_{\overline{M}/(x_{i+2}, \dots, x_{d-1})\overline{M}}).$$

It follows by inductive hypothesis that $x_1, \dots, x_{d-1}$ is a dd-sequence of $\overline{M}$. Therefore we have only to prove that $\underline{x}$ is a strong d-sequence of M . Since $x_1, \dots, x_{d-1}$ is a strong d-sequence of $\overline{M}$,

$$(x_d^{n_d}, x_1^{n_1}, \dots, x_i^{n_i})M : x_j^{n_j} = (x_d^{n_d}, x_1^{n_1}, \dots, x_i^{n_i})M : x_{i+1}^{n_{i+1}} x_j^{n_j}$$

for all $0 \leq i < j \leq d - 1$. Hence, by Krull's Intersection Theorem,

$$\begin{aligned}(x_1^{n_1}, \dots, x_i^{n_i})M : x_j^{n_j} &= \bigcap_{n_d} (x_d^{n_d}, x_1^{n_1}, \dots, x_i^{n_i})M : x_j^{n_j} \\ &= \bigcap_{n_d} (x_d^{n_d}, x_1^{n_1}, \dots, x_i^{n_i})M : x_{i+1}^{n_{i+1}} x_j^{n_j} \\ &= (x_1^{n_1}, \dots, x_i^{n_i})M : x_{i+1}^{n_{i+1}} x_j^{n_j}.\end{aligned}$$

So in order to show that $\underline{x}$ is a strong d-sequence of M . It remains to check that

$$(x_1^{n_1}, \dots, x_i^{n_i})M : x_d^{n_d} = (x_1^{n_1}, \dots, x_i^{n_i})M : x_{i+1}^{n_{i+1}} x_d^{n_d}, \text{ for all } 0 \leq i < d.$$

In fact, applying Lemma 2.1 to the sequence $x_d^{n_d}, x_{i+1}^{n_{i+1}}, x_1^{n_1}, \dots, \widehat{x_{i+1}^{n_{i+1}}}, \dots, x_{d-1}^{n_{d-1}}$, we obtain for all $i = 0, \dots, d-1$ that

$$\begin{aligned} \chi_1(\underline{x}(\underline{n}); M) &= \sum_{\substack{j=0 \\ j \neq i}}^{d-2} e(x_d^{n_d}, x_{i+1}^{n_{i+1}}, x_1^{n_1}, \dots, x_j^{n_j}; (0 : x_{j+1}^{n_{j+1}})_{M/(x_{j+2}^{n_{j+2}}, \dots, \widehat{x_{i+1}^{n_{i+1}}}, \dots, x_{d-1}^{n_{d-1}})M}) \\ &\quad + n_d e(x_d; (0 : x_{i+1}^{n_{i+1}})_{M/(x_1^{n_1}, \dots, \widehat{x_{i+1}^{n_{i+1}}}, \dots, x_{d-1}^{n_{d-1}})M}) \\ &\quad + \ell((0 : x_d^{n_d})_{M/(x_{i+1}^{n_{i+1}}, x_1^{n_1}, \dots, x_{d-1}^{n_{d-1}})M}). \end{aligned}$$

Moreover, since $\chi_1(\underline{x}(\underline{n}); M)$ does not depend on n_d , we deduce that

$$e(x_d; (0 : x_{i+1}^{n_{i+1}})_{M/(x_1^{n_1}, \dots, \widehat{x_{i+1}^{n_{i+1}}}, \dots, x_{d-1}^{n_{d-1}})M}) = 0.$$

Thus there exists a positive integer n such that

$$x_d^n (0 : x_{i+1}^{n_{i+1}})_{M/(x_1^{n_1}, \dots, \widehat{x_{i+1}^{n_{i+1}}}, \dots, x_{d-1}^{n_{d-1}})M} = 0.$$

So

$$(0 : x_{i+1}^{n_{i+1}})_{M/(x_1^{n_1}, \dots, \widehat{x_{i+1}^{n_{i+1}}}, \dots, x_{d-1}^{n_{d-1}})M} \subseteq (0 : x_d^n)_{M/(x_1^{n_1}, \dots, \widehat{x_{i+1}^{n_{i+1}}}, \dots, x_{d-1}^{n_{d-1}})M}.$$

On the other hand, since $\chi_1(\underline{x}(\underline{n}); M)$ is a polynomial, it follows by [4, Theorem 1] that

$$(0 : x_d^{n+n_d})_{M/(x_1^{n_1}, \dots, \widehat{x_{i+1}^{n_{i+1}}}, \dots, x_{d-1}^{n_{d-1}})M} = (0 : x_d^{n_d})_{M/(x_1^{n_1}, \dots, \widehat{x_{i+1}^{n_{i+1}}}, \dots, x_{d-1}^{n_{d-1}})M}$$

for all n_d . Therefore

$$\begin{aligned} (0 : x_{i+1}^{n_{i+1}} x_d^{n_d})_{M/(x_1^{n_1}, \dots, \widehat{x_{i+1}^{n_{i+1}}}, \dots, x_{d-1}^{n_{d-1}})M} &\subseteq (0 : x_d^{n+n_d})_{M/(x_1^{n_1}, \dots, \widehat{x_{i+1}^{n_{i+1}}}, \dots, x_{d-1}^{n_{d-1}})M} \\ &= (0 : x_d^{n_d})_{M/(x_1^{n_1}, \dots, \widehat{x_{i+1}^{n_{i+1}}}, \dots, x_{d-1}^{n_{d-1}})M}. \end{aligned}$$

Thus

$$(x_1^{n_1}, \dots, \widehat{x_{i+1}^{n_{i+1}}}, \dots, x_{d-1}^{n_{d-1}})M : x_d^{n_d} = (x_1^{n_1}, \dots, \widehat{x_{i+1}^{n_{i+1}}}, \dots, x_{d-1}^{n_{d-1}})M : x_{i+1}^{n_{i+1}} x_d^{n_d}.$$

Now, applying Krull's Intersection Theorem again we get

$$\begin{aligned} (x_1^{n_1}, \dots, x_i^{n_i})M : x_d^{n_d} &= \bigcap_{n_{i+2}, \dots, n_{d-1}} (x_1^{n_1}, \dots, \widehat{x_{i+1}^{n_{i+1}}}, \dots, x_{d-1}^{n_{d-1}})M : x_d^{n_d} \\ &= \bigcap_{n_{i+2}, \dots, n_{d-1}} (x_1^{n_1}, \dots, \widehat{x_{i+1}^{n_{i+1}}}, \dots, x_{d-1}^{n_{d-1}})M : x_{i+1}^{n_{i+1}} x_d^{n_d} \\ &= (x_1^{n_1}, \dots, x_i^{n_i})M : x_{i+1}^{n_{i+1}} x_d^{n_d} \end{aligned}$$

for all $i = 0, \dots, d-1$. So $\underline{x}$ is a strong d-sequence and the proof of Theorem 1.2 is complete. $\square$

As an application of Theorem 1.2 we get the following useful criterion to check whether a system of parameters is a dd-sequence.

Corollary 3.6. *A system of parameters $\underline{x}$ of M is a dd-sequence of M if and only if there exist integers $a_0, a_1, \dots, a_{d-1}$ such that*

$$\chi_1(\underline{x}(\underline{n}); M) = \sum_{i=0}^{d-1} n_1 \dots n_i a_i.$$

Moreover, in this case it holds

$$a_i = e(x_1, \dots, x_i; (0 : x_{i+1})_{M/(x_{i+2}, \dots, x_d)M})$$

for $i = 0, 1, \dots, p_1(M)$ and $a_{p_1(M)+1} = \dots = a_{d-1} = 0$.

Proof. The necessary condition follows immediately Theorem 1.2. We proceed by induction on d that

$$a_i = e(x_1, \dots, x_i; (0 : x_{i+1})_{M/(x_{i+2}, \dots, x_d)M})$$

for all $i = 0, 1, \dots, d-1$, and the corollary follows by Theorem 1.2. In fact, the case $d = 1$ is obvious. Suppose that $d > 1$. We set $M_d = M/x_d M$ and $\underline{x}'(\underline{n}') = (x_1^{n_1}, \dots, x_{d-1}^{n_{d-1}})$. Then

$$\begin{aligned} \chi_1(\underline{x}'(\underline{n}'); M_d) &= \chi_1(\underline{x}(\underline{n}); M) - e(x_1^{n_1}, \dots, x_{d-1}^{n_{d-1}}; 0 :_M x_d) \\ &= \sum_{i=0}^{d-2} n_1 \dots n_i a_i + n_1 \dots n_{d-1} (a_{d-1} - e(x_1, \dots, x_{d-1}; 0 :_M x_d)). \end{aligned}$$

Since the function $\chi_1(x_1^{n_1}, \dots, x_{d-1}^{n_{d-1}}; M_d)$ is increasing and bounded above by a polynomial in $n_1, \dots, n_{d-1}$ of degree $d-2$, it implies that

$$a_{d-1} = e(x_1, \dots, x_{d-1}; 0 :_M x_d).$$

Therefore

$$\chi_1(x_1^{n_1}, \dots, x_{d-1}^{n_{d-1}}; M_d) = \sum_{i=0}^{d-2} n_1 \dots n_i a_i.$$

So we get by the inductive hypothesis that

$$a_i = e(x_1, \dots, x_i; (0 : x_{i+1})_{M/(x_{i+2}, \dots, x_d)M}),$$

for $i = 0, \dots, d-2$. $\square$

The following corollary is an immediate consequence of Theorem 1.2.

Corollary 3.7. *Let $\underline{x} = (x_1, \dots, x_d)$ be a system of parameters of M and $k = p_1(M)$. Suppose that $\underline{x}$ is a dd-sequence of M . Then, for any permutation π of the set $\{k+1, \dots, d\}$, $x_1, \dots, x_k, x_{\pi(k+1)}, \dots, x_{\pi(d)}$ is also a dd-sequence.*

Remark 3.8. (i) Recall that an R -module M is said to be a generalized Cohen-Macaulay module, if $\sup_{\underline{x}} \{\chi_1(\underline{x}; M)\} = I(M) < \infty$, where $\underline{x}$ runs through all

systems of parameters of M . It was shown in [24] that M is a generalized Cohen-Macaulay module if and only if there exists a system of parameters $\underline{x}$ of M such that $\chi_1(\underline{x}; M) = I(M)$. That system of parameters was called a standard system of parameters of M and plays an important role in the theory of generalized Cohen-Macaulay modules. Suppose now that M is a generalized Cohen-Macaulay module and $\underline{x}$ a system of parameters of M . Then, it follows from Theorem 1.2 that $\underline{x}$ is a standard system of parameters if and only if $\underline{x}$ is a dd-sequence. Moreover, an R -module M is called a Buchsbaum module if every system of parameters of M is a standard system of parameters. Therefore we get the following characterization of Buchsbaum modules by dd-sequences as follows: M is a Buchsbaum module if and only if every system of parameters of M is a dd-sequence.

(ii) Let $\mathfrak{a}_i(M) = \text{Ann}_R(H_{\mathfrak{m}}^i(M))$ be the annihilator ideal of the i -th local cohomology module $H_{\mathfrak{m}}^i(M)$ of M and $\mathfrak{a}(M) = \mathfrak{a}_0(M) \dots \mathfrak{a}_{d-1}(M)$, where $d = \dim M$. Then, a system of parameters $\underline{x}$ of M was called a *p-standard system of parameters* if $x_i \in \mathfrak{a}(M/(x_{i+1}, \dots, x_d)M)$ for $i = 1, \dots, d$. This kind of system of parameters was introduced and investigated in [5, 6, 7] and has been used for solving the problem of Macaulayfication by recent works of T. Kawasaki [16, 17]. It was proved in [7] that if $\underline{x}$ is a p-standard system of parameters of M , then

$$\chi_1(\underline{x}(\underline{n}); M) = \sum_{i=0}^{p_1(M)} n_1 \dots n_i e(x_1, \dots, x_i; (0 : x_{i+1})_{M/(x_{i+2}, \dots, x_d)M}).$$

Therefore any p-standard system of parameters is a dd-sequence. Unfortunately, the converse is wrong. It means that a system of parameters, which is a dd-sequence, in general is not a p-standard system of parameters (see Example 3.11). However, the following consequence gives us close relations between them.

Corollary 3.9. *Let $\underline{x} = (x_1, \dots, x_d)$ be a system of parameters of M . If $\underline{x}$ is a p-standard system of parameters then it is also a dd-sequence. Conversely, suppose that $\underline{x}$ is a dd-sequence. Then $x_1^{n_1}, \dots, x_d^{n_d}$ is a p-standard system of parameters for all $n_i \geq i$, $i = 1, \dots, d$.*

Proof. The first statement is already shown in the remark above. Since a dd-sequence is a strong d-sequence, the second one is just an immediate consequence of Lemma 2.9 in [7], which says that

$$x_j H_{\mathfrak{m}}^i(M/(x_1^{n_1}, \dots, x_h^{n_h})M) = 0$$

for all $j = 1, \dots, d$, $h + i < j$ and $n_1, \dots, n_h > 0$, provided the system of parameters $\underline{x} = (x_1, \dots, x_d)$ is a strong d-sequence. It should be noticed that

this result was stated in [7] for p-standard systems of parameters, but in its proof one needs only the condition that the system of parameters $\underline{x}$ is a strong d-sequence. $\square$

As we have seen, a p-standard system of parameters is a dd-sequence and a dd-sequence is always a strong d-sequence. However, the converse statements are wrong in general. Below we present some examples to clarify these three notions.

Example 3.10. Let $R = k[[X, Y]]$ be the ring of all formal power series of two indeterminates X, Y over a field k with the maximal ideal $\mathfrak{m}$. Consider $M = \mathfrak{m}^2$ as an R -module. Then $\dim(M) = 2$ and X, Y^2 form a system of parameters of M . It is easy to check that $0 :_M X^m Y^{2n} = 0 = 0 :_M Y^{2n}$ and $X^m M : Y^{2n} = X^m M : Y^2$ for all positive integers m, n . Therefore X, Y^2 is a strong d-sequence of M . On the other hand, we can show that

$$Y^2 M : X^m = \begin{cases} (XY^2, Y^3), & \text{if } m = 1, \\ (Y^2), & \text{if } m > 1. \end{cases}$$

Hence X, Y^2 is not a dd-sequence of M . Besides, we can also compute the 1-st Euler-Poincaré characteristic and get

$$\chi_1(X^m, Y^{2n}; M) = \begin{cases} 2, & \text{if } m = 1, \\ 3, & \text{if } m > 1. \end{cases}$$

Example 3.11. Let $R = k[[X_1, \dots, X_{d+1}]]$, ($d > 1$) be the ring of all formal power series of $(d + 1)$ -indeterminates $X_1, \dots, X_{d+1}$ over a field k with the maximal ideal $\mathfrak{m} = (X_1, \dots, X_{d+1})$. Let M denote the R -module R/I where $I = (X_{d+1}^{d+1}, X_1 X_{d+1}^d, X_2 X_{d+1}^{d-1}, \dots, X_d X_{d+1})$. It is easy to see that $\dim(M) = d$ and $X_1, \dots, X_d$ is a system of parameters of M . Then by a simple computation we get for all $n_1, \dots, n_d \in \mathbb{N}^d$,

$$\ell(M/(X_1^{n_1}, \dots, X_d^{n_d})M) = \sum_{i=0}^d n_1 \dots n_i,$$

where we set $n_1 \dots n_i = 1$ if $i = 0$. Hence

$$\chi_1(X_1^{n_1}, \dots, X_d^{n_d}; M) = \sum_{i=0}^{d-1} n_1 \dots n_i.$$

Therefore the system of parameters $X_1, \dots, X_d$ is a dd-sequence of M with $p_1(M) = d - 1$. On the other hand, we can check that

$$H_{\mathfrak{m}}^0(M) \cong (X_{d+1}^d, X_2 X_{d+1}^{d-1}, \dots, X_d X_{d+1})/I.$$

Hence $\mathfrak{a}_0(M) = \text{Ann}_R(H_{\mathfrak{m}}^0(M)) = \mathfrak{m}$. Moreover, since R is a regular local ring, we have $\dim(R/\mathfrak{a}(M)) = p_1(M) = d - 1 > 0$ by [5, Theorem 1.2]. It follows that there exists at least an $i \in \{1, \dots, d - 1\}$ such that $\mathfrak{a}_i(M) \subseteq \mathfrak{m}$. Therefore $\mathfrak{a}(M) \subseteq \mathfrak{a}_0(M)\mathfrak{a}_i(M) \subseteq \mathfrak{m}^2$. Since $X_d \notin \mathfrak{m}^2$, $X_d \notin \mathfrak{a}(M)$. Thus $X_1, \dots, X_d$ is not a p-standard system of parameters of M .

4 Partial Euler-Poincaré characteristics with respect to a dd-sequence

Keep all notations in the previous sections. We begin this Section with the following key lemma.

Lemma 4.1. *Suppose that $\underline{x} = (x_1, \dots, x_s)$ is a dd-sequence of M . Then*

$$(0 : x_j)_{H_k(x_1, \dots, x_{i-1}, x_{j+1}, \dots, x_s; M)} = (0 : x_i x_j)_{H_k(x_1, \dots, x_{i-1}, x_{j+1}, \dots, x_s; M)}$$

for all $1 \leq i \leq j \leq s$ and $0 \leq k \leq s$.

Proof. Since the lemma is trivial for the case $s = 1$, we assume that $s \geq 2$. Moreover, since $x_1, \dots, x_{i-1}, x_i, x_j, x_{j+1}, \dots, x_s$ is a dd-sequence of M by the hypothesis and Corollary 3.5, it suffices to prove the lemma for only two cases $j = i + 1$ and $j = i$. We will do it for the case $j = i + 1$ and the proof for the other case is quite similar. In fact, we need to show that

$$(0 : x_{i+1})_{H_k(x_1, \dots, x_{i-1}, x_{i+2}, \dots, x_s; M)} = (0 : x_i x_{i+1})_{H_k(x_1, \dots, x_{i-1}, x_{i+2}, \dots, x_s; M)}. \quad (*)$$

Set $\underline{y} = (y_1, \dots, y_s)$, where

$$y_j = \begin{cases} x_j & \text{if } j < i, \\ x_{j+2} & \text{if } i \leq j \leq s-2, \\ x_i & \text{if } j = s-1, \\ x_{i+1} & \text{if } j = s. \end{cases}$$

Then the above equality $(*)$ can be rewritten as follows

$$(0 : y_s)_{H_k(y_1, \dots, y_{s-2}; M)} = (0 : y_{s-1} y_s)_{H_k(y_1, \dots, y_{s-2}; M)}. \quad (**)$$

Denote by φ_k the k -th differential of the Koszul complex $K(y_1, \dots, y_{s-2}; M)$. Since $H_k(y_1, \dots, y_{s-2}; M) = \ker(\varphi_k)/\text{im}(\varphi_{k+1})$, the equality $(**)$ and hence the lemma are proved if we can verify the following claim.

Claim. *With all notations above it holds*

$$\text{im}(\varphi_{k+1}) \cdot_{\ker(\varphi_k)} y_s = \text{im}(\varphi_{k+1}) \cdot_{\ker(\varphi_k)} y_s y_{s-1}.$$

for all $s \geq 2$ and $k = 0, 1, \dots, s$.

Proof of the Claim. Recall that any element $a \in K_k(y_1, \dots, y_{s-2}; M) \cong M^{\binom{s-2}{k}}$ can be expressed as $a = (a_{i_1, \dots, i_k})_{1 \leq i_1 < i_2 < \dots < i_k \leq s-2} \in M^{\binom{s-2}{k}}$. Extend the $a_{i_1, \dots, i_k}$'s to an alternating function of the indices, i. e. such that $a_{\dots, i, \dots, i, \dots} = 0$ and $a_{\dots, i, \dots, j, \dots} = -a_{\dots, j, \dots, i, \dots}$ for all $i \neq j$. For convenience, we say in this case that the elements $a_{i_1, \dots, i_k} \in M$ are of alternating indices. Then we have

$$\ker(\varphi_k) = \{(a_{i_1, \dots, i_k})_{1 \leq i_1 < i_2 < \dots < i_k \leq s-2} \in M^{\binom{s-2}{k}} \mid \sum_{j=1}^{s-2} y_j a_{j, i_1, \dots, i_{k-1}} = 0\},$$

$$\begin{aligned} \text{im}(\varphi_{k+1}) &= \{(b_{i_1, \dots, i_k})_{1 \leq i_1 < i_2 < \dots < i_k \leq s-2} \in M^{\binom{s-2}{k}} | \\ &\exists (a_{i_1, \dots, i_{k+1}})_{1 \leq i_1 < \dots < i_{k+1} \leq s-2} \in M^{\binom{s-2}{k+1}} : b_{i_1, \dots, i_k} = \sum_{j=1}^{s-2} y_j a_{j, i_1, \dots, i_k}\}. \end{aligned}$$

We prove the claim by induction on s . The case $s = 2$ is straightforward. Now assume that $s > 2$. Let $(a_{i_1, \dots, i_k})_{1 \leq i_1 < \dots < i_k \leq s-2}$ be an arbitrary element of $\text{im}(\varphi_{k+1}) :_{\ker(\varphi_k)} y_{s-1} y_s$. Then, it follows from the above descriptions of $\ker(\varphi_k)$ and $\text{im}(\varphi_{k+1})$ that

$$(a_{i_1, \dots, i_k})_{1 \leq i_1 < \dots < i_k \leq s-2} \in \text{im}(\varphi_{k+1}) :_{\ker(\varphi_k)} y_s$$

if there exist elements $b_{j, i_1, \dots, i_k} \in M$ of alternating indices such that

$$y_s a_{i_1, \dots, i_k} = \sum_{j=1}^{s-2} y_j b_{j, i_1, \dots, i_k}. \quad (***)$$

Therefore the claim is proved if we can show the existence of $b_{j, i_1, \dots, i_k}$ and the equality (***). To do it, we consider the following cases.

Case 1: $i_1 > 1$. Put $\overline{M} = M/y_1 M$ and $\overline{a}_{i_1, \dots, i_k}$ the image of $a_{i_1, \dots, i_k}$ in $\overline{M}$. Denote by φ'_k the k -th differential of the Koszul complex $K(y_2, \dots, y_{s-2}; \overline{M})$. Then it follows from Proposition 3.4 and the inductive hypothesis that

$$(\overline{a}_{i_1, \dots, i_k})_{i_1 < \dots < i_k \leq s-2} \in \text{im}(\varphi'_{k+1}) :_{\ker(\varphi'_k)} y_{s-1} y_s = \text{im}(\varphi'_{k+1}) :_{\ker(\varphi'_k)} y_s.$$

So there exist elements $b_{j, i_1, \dots, i_k} \in M$ of alternating indices such that

$$y_s a_{i_1, \dots, i_k} = \sum_{j=1}^{s-2} y_j b_{j, i_1, \dots, i_k}.$$

Case 2: $i_1 = 1$. We have to prove in this case that

$$y_s a_{1, i_2, \dots, i_k} = \sum_{j=2}^{s-2} y_j b_{j, 1, i_2, \dots, i_k},$$

where $b_{j, 1, \dots, i_k}$'s are of alternating indices and just determined in the case 1. Since $x_1, \dots, x_s$ is a dd-sequence of M , we can apply Proposition 3.4 several times to imply that the sequence x_i, x_{i+1} is a dd-sequence of the module $M/(x_1, \dots, x_{i-1}, x_{i+2}, \dots, x_s)M$. Therefore y_{s-1}, y_s is a d-sequence with respect to $M/(y_1, \dots, y_{s-2})M$. Hence, from the hypothesis

$$y_{s-1} y_s a_{1, i_2, \dots, i_k} = \sum_{j=2}^{s-2} y_j a_{j, 1, i_2, \dots, i_k},$$

it follows that

$$a_{1, i_2, \dots, i_k} \in \left(\sum_{j=2}^{s-2} y_j R \right) M : y_{s-1} y_s = \left(\sum_{j=2}^{s-2} y_j R \right) M : y_s.$$

Let t be a positive number such that

$$y_s a_{1,i_2,\dots,i_k} = \sum_{j=2}^{t-1} y_j b_{j,1,i_2,\dots,i_k} + \sum_{j=t}^{s-2} y_j c_j,$$

where $c_t, \dots, c_{s-2} \in M$. Then $t \geq 2$ and the first summand of the right term is 0 if $t = 2$. We put $I = (y_{t+1}, \dots, y_{s-2})R$, $M' = M/IM$ and denote by a' the image of the element $a \in M$ in M' . We get

$$y_s a'_{1,i_2,\dots,i_k} = \sum_{j=2}^{t-1} y_j b'_{j,1,i_2,\dots,i_k} + y_t c'_t$$

and by the case 1 that

$$y_s a'_{i_1,\dots,i_k} = \sum_{j=1}^t y_j b'_{j,i_1,\dots,i_k}, \quad (i_1 > 1).$$

Therefore (note that $b_{i,j,i_2,\dots,i_k}$ are of alternating indices)

$$\begin{aligned} y_s \sum_{j=1}^t y_j a'_{j,i_2,\dots,i_k} &= \sum_{j=1}^t y_j (y_s a'_{j,i_2,\dots,i_k}) \\ &= \sum_{j=2}^t y_j (y_s a'_{j,i_2,\dots,i_k}) + y_1 (y_s a'_{1,i_2,\dots,i_k}) \\ &= \sum_{j=2}^t \sum_{i=1}^t y_j y_i b'_{i,j,i_2,\dots,i_k} + \sum_{j=2}^{t-1} y_1 y_j b'_{j,1,i_2,\dots,i_k} + y_1 y_t c'_t \\ &= y_1 y_t (b'_{1,t,i_2,\dots,i_k} + c'_t) \\ &= y_1 y_t (c'_t - b'_{t,1,i_2,\dots,i_k}). \end{aligned}$$

On the other hand, since $(a_{j,i_2,\dots,i_k})_{j,i_2,\dots,i_k} \in \ker(\varphi_k)$,

$$\sum_{j=1}^{s-2} y_j a_{j,i_2,\dots,i_k} = 0, \quad \text{thus} \quad \sum_{j=1}^t y_j a'_{j,i_2,\dots,i_k} = 0.$$

Hence

$$0 = y_s \sum_{j=1}^t y_j a'_{j,i_2,\dots,i_k} = y_1 y_t (c'_t - b'_{t,1,i_2,\dots,i_k}).$$

Moreover, since $\underline{x}$ is a dd-sequence of M , it is a strong d-sequence. We can show by applying Krull's Intersection Theorem that $(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_{s-2}) = (y_1, \dots, y_{s-2})$ is also a strong d-sequence. Hence $y_1, \dots, y_t$ is a d-sequence of M' . So we have $c_t - b_{t,1,i_2,\dots,i_k} \in IM :_M y_1 y_t = IM :_M y_t$. Therefore $y_t (c_t - b_{t,1,i_2,\dots,i_k}) \in IM$. It follows that there exist $d_{t+1}, \dots, d_{s-2} \in M$ such that

$$y_s a_{1,i_2,\dots,i_k} = \sum_{j=2}^t y_j b_{j,1,i_2,\dots,i_k} + \sum_{j=t+1}^{s-2} y_j d_j.$$

If $t < s - 2$, we can repeat the above process. Finally, after at most $(s - 2 - t)$ -steps, we get

$$y_s a_{1,i_2,\dots,i_k} = \sum_{j=2}^{s-2} y_j b_{j,1,i_2,\dots,i_k}$$

as required and the proof of the lemma is complete. $\square$

The following lemma proved in [9] is needed for the proofs of the theorem 1.3 and its consequences.

Lemma 4.2. [9, Lemma 3.1] *Let $\underline{x} = (x_1, \dots, x_d)$ be a system of parameters of M . If $\underline{x}$ is a strong d -sequence, then for $0 < i \leq j \leq d$ the length of the Koszul homology module $H_i(x_1^{n_1}, \dots, x_j^{n_j}; M)$ is finite and given by*

$$\ell(H_i(x_1^{n_1}, \dots, x_j^{n_j}; M)) = \sum_{t=0}^{j-i} \binom{j-t-1}{i-1} \ell(H_{\mathfrak{m}}^0(M/(x_1^{n_1}, \dots, x_t^{n_t})M)).$$

Therefore the function $\ell(H_i(x_1^{n_1}, \dots, x_j^{n_j}; M))$ depends only on $n_1, \dots, n_{j-i}$.

Now we are able to prove Theorem 1.3.

Proof of Theorem 1.3. Since $\underline{x}$ is a dd-sequence, $\chi_{d-k}(\underline{x}(\underline{n}); M)$ is a polynomial in $\underline{n}$ by Theorem 1.1 and Lemma 4.1. Then $\deg \chi_{d-k}(\underline{x}(\underline{n}); M) = p_{d-k}(M) \leq k$ by Remark 2.3. Therefore the theorem is proved if we can show that

$$\chi_{d-k}(\underline{x}(\underline{n}); M) = \sum_{i=0}^k n_1 \dots n_i e(x_1, \dots, x_i; (0 : x_{i+1})_{H_{d-k-1}(x_{i+2}, \dots, x_d; M)}) \quad (\#)$$

for all $k = 0, 1, \dots, d - 1$. We will do this by the recursive method. Denote by $\underline{x}_{k+1}$ the sequence $x_{k+1}, \dots, x_d$. Then by virtue of Lemma 4.2 and applying Lemma 2.1 to the sequence $x_1^{n_1}, \dots, x_k^{n_k}, x_{k+1}, \dots, x_d$ we get

$$\begin{aligned} \chi_{d-k}(\underline{x}(\underline{n}); M) &= \chi_{d-k}(x_1^{n_1}, \dots, x_k^{n_k}, \underline{x}_{k+1}; M) \\ &= n_1 \dots n_k e(x_1, \dots, x_k; (0 : x_{k+1})_{H_{d-k-1}(\underline{x}_{k+2}; M)}) \\ &\quad + \sum_{i=0}^{k-1} n_1 \dots n_i e(x_1, \dots, x_i; (0 : x_{i+1}^{n_{i+1}})_{H_{d-k-1}(x_{i+2}^{n_{i+2}}, \dots, x_k^{n_k}, \underline{x}_{k+1}; M)}). \end{aligned}$$

Put

$$\begin{aligned} \phi_1(n_1, \dots, n_k) &= \chi_{d-k}(x_1^{n_1}, \dots, x_k^{n_k}, \underline{x}_{k+1}; M) \\ &\quad - n_1 \dots n_k e(x_1, \dots, x_k; (0 : x_{k+1})_{H_{d-k-1}(\underline{x}_{k+2}; M)}) \\ &= \sum_{i=0}^{k-1} n_1 \dots n_i e(x_1, \dots, x_i; (0 : x_{i+1}^{n_{i+1}})_{H_{d-k-1}(x_{i+2}^{n_{i+2}}, \dots, x_k^{n_k}, \underline{x}_{k+1}; M)}). \end{aligned}$$

First, we claim that the function $\phi_1(n_1, \dots, n_k)$ is independent of n_k . In fact, since the function $\chi_{d-k}(\underline{x}(\underline{n}); M)$ does not depend on the order of $x_1, \dots, x_d$,

using again Lemma 2.1 to the sequence $x_k^{n_k}, x_{k-1}^{n_{k-1}}, \dots, x_1^{n_1}, x_{k+1}, \dots, x_d$ we obtain

$$\begin{aligned} \chi_{d-k}(\underline{x}(\underline{n})); M) &= n_1 \dots n_k e(x_1, \dots, x_k; (0 : x_{k+1})_{H_{d-k-1}(\underline{x}_{k+2}; M)}) \\ &\quad + \sum_{i=2}^{k+1} n_i \dots n_k e(x_i, \dots, x_k; (0 : x_{i-1}^{n_{i-1}})_{H_{d-k-1}(x_1^{n_1}, \dots, x_{i-2}^{n_{i-2}}, \underline{x}_{k+1}; M)}). \end{aligned}$$

Hence

$$\begin{aligned} \phi_1(n_1, \dots, n_k) &= \sum_{i=2}^k n_i \dots n_k e(x_i, \dots, x_k; (0 : x_{i-1}^{n_{i-1}})_{H_{d-k-1}(x_1^{n_1}, \dots, x_{i-2}^{n_{i-2}}, \underline{x}_{k+1}; M)}) \\ &\quad + \ell((0 : x_k^{n_k})_{H_{d-k-1}(x_1^{n_1}, \dots, x_{k-1}^{n_{k-1}}, \underline{x}_{k+1}; M)}). \end{aligned}$$

On the other hand, it follows by Lemma 4.1 that

$$\begin{aligned} (0 : x_{i-1}^{n_{i-1}})_{H_{d-k-1}(x_1^{n_1}, \dots, x_{i-2}^{n_{i-2}}, \underline{x}_{k+1}; M)} &\subseteq (0 : x_{i-1}^{n_{i-1}} x_k)_{H_{d-k-1}(x_1^{n_1}, \dots, x_{i-2}^{n_{i-2}}, \underline{x}_{k+1}; M)} \\ &= (0 : x_k)_{H_{d-k-1}(x_1^{n_1}, \dots, x_{i-2}^{n_{i-2}}, \underline{x}_{k+1}; M)} \end{aligned}$$

for all $2 \leq i \leq k$. Thus $x_k(0 : x_{i-1}^{n_{i-1}})_{H_{d-k-1}(x_1^{n_1}, \dots, x_{i-2}^{n_{i-2}}, \underline{x}_{k+1}; M)} = 0$. So $e(x_i, \dots, x_k; (0 : x_{i-1}^{n_{i-1}})_{H_{d-k-1}(x_1^{n_1}, \dots, x_{i-2}^{n_{i-2}}, \underline{x}_{k+1}; M)}) = 0$. Therefore, by Lemma 4.1,

$$\begin{aligned} \phi_1(n_1, \dots, n_k) &= \ell((0 : x_k^{n_k})_{H_{d-k-1}(x_1^{n_1}, \dots, x_{k-1}^{n_{k-1}}, \underline{x}_{k+1}; M)}) \\ &= \ell((0 : x_k)_{H_{d-k-1}(x_1^{n_1}, \dots, x_{k-1}^{n_{k-1}}, \underline{x}_{k+1}; M)}) \end{aligned}$$

is independent of n_k . Thus the formula (#) is proved if $k = 1$. Assume that $k \geq 2$. Then

$$\begin{aligned} \chi_{d-k}(\underline{x}(\underline{n}); M) &= n_1 \dots n_k e(x_1, \dots, x_k; (0 : x_{k+1})_{H_{d-k-1}(\underline{x}_{k+2}; M)}) + \phi_1(n_1, \dots, n_{k-1}, 1) \\ &= n_1 \dots n_k e(x_1, \dots, x_k; (0 : x_{k+1})_{H_{d-k-1}(\underline{x}_{k+2}; M)}) \\ &\quad + n_1 \dots n_{k-1} e(x_1, \dots, x_{k-1}; (0 : x_k)_{H_{d-k-1}(\underline{x}_{k+1}; M)}) \\ &\quad + \sum_{i=0}^{k-2} n_1 \dots n_i e(x_1, \dots, x_i; (0 : x_{i+1}^{n_{i+1}})_{H_{d-k-1}(x_{i+2}^{n_{i+2}}, \dots, x_{k-1}^{n_{k-1}}, J_k; M)}). \end{aligned}$$

Next, we set

$$\begin{aligned} \phi_2(n_1, \dots, n_{k-1}) &= \chi_{d-k}(\underline{x}(\underline{n}); M) - n_1 \dots n_k e(x_1, \dots, x_k; (0 : x_{k+1})_{H_{d-k-1}(\underline{x}_{k+2}; M)}) \\ &\quad - n_1 \dots n_{k-1} e(x_1, \dots, x_{k-1}; (0 : x_k)_{H_{d-k-1}(\underline{x}_{k+1}; M)}) \\ &= \sum_{i=0}^{k-2} n_1 \dots n_i e(x_1, \dots, x_i; (0 : x_{i+1}^{n_{i+1}})_{H_{d-k-1}(x_{i+2}^{n_{i+2}}, \dots, x_{k-1}^{n_{k-1}}, J_k; M)}). \end{aligned}$$

With the same method as used above, we can verify that $\phi_2(n_1, \dots, n_{k-1})$ is independent of n_{k-1} . Continuing this procedure, after k - steps, we obtain

$$\chi_{d-k}(\underline{x}(\underline{n}); M) = \sum_{i=0}^k n_1 \dots n_i e(x_1, \dots, x_i; (0 : x_{i+1})_{H_{d-k-1}(x_{i+2}, \dots, x_d; M)})$$

as required. $\square$

Corollary 4.3. *Let $\underline{x} = (x_1, \dots, x_d)$ be a system of parameters of M . If $\underline{x}$ is a dd -sequence, then $\ell(H_k(x_1^{n_1}, \dots, x_d^{n_d}; M))$ is a polynomial in $n_1, \dots, n_{p_k(M)}$ of degree $p_k(M)$ for all $k = 0, 1, \dots, d$.*

Proof. Since

$$\chi_k(\underline{x}(\underline{n}); M) + \chi_{k+1}(\underline{x}(\underline{n}); M) = \ell(H_k(\underline{x}(\underline{n}); M)),$$

the statement follows from Theorem 1.3 and the fact that

$$\deg(\chi_k(\underline{x}(\underline{n}); M)) = p_k(M) \geq p_{k+1}(M).$$

$\square$

5 Local Cohomology Modules

Recall that a system of parameters $\underline{x} = (x_1, \dots, x_d)$ of M is said to be a standard system of parameters if $\chi_1(\underline{x}(\underline{n}); M)$ is a constant for all $n_1, \dots, n_d > 0$ and that the module M is a generalized Cohen-Macaulay module if there exists a standard system of parameters on M , i. e. $p_1(M) \leq 0$. It is well-known that M is a generalized Cohen-Macaulay module if and only if $\ell(H_{\mathfrak{m}}^i(M)) < \infty$ for all $i = 1, \dots, d-1$.

Lemma 5.1. *Let $\underline{x} = (x_1, \dots, x_d)$ be a standard system of parameters of a generalized Cohen-Macaulay module M . Then*

$$\ell(H_{\mathfrak{m}}^i(M)) = \sum_{j=0}^i (-1)^{i-j} \binom{i}{j} \ell(H_{\mathfrak{m}}^0(M/(x_1, \dots, x_j)M))$$

for all $i < d$.

Proof. We proceed by induction on d and i . The lemma is trivial if $d = 0, 1$ or $i = 0$. Assume that $d > 1$ and $i > 0$. Since $\underline{x}$ is a standard system of parameters of M , $(x_1, \dots, x_d)H_{\mathfrak{m}}^i(M) = 0$ for all $i < d$. Hence we have the following exact sequences

$$0 \longrightarrow H_{\mathfrak{m}}^{i-1}(M) \longrightarrow H_{\mathfrak{m}}^{i-1}(M/x_1M) \longrightarrow H_{\mathfrak{m}}^i(M) \longrightarrow 0$$

for all $0 < i < d$. Therefore

$$\ell(H_{\mathfrak{m}}^i(M)) = \ell(H_{\mathfrak{m}}^{i-1}(M/x_1M)) - \ell(H_{\mathfrak{m}}^{i-1}(M))$$

and the lemma follows easily from the inductive hypotheses. $\square$

A consequence of Theorem 1.2 is that if $p_1(M) > 0$ and the system of parameters $\underline{x} = (x_1, \dots, x_d)$ is a dd-sequence of M then $p_1(M/x_1M) = p_1(M) - 1$. Therefore, if $k \geq p_1(M)$, $p_1(M/(x_1, \dots, x_k)M) \leq 0$. So $M/(x_1, \dots, x_k)M$ is a generalized Cohen-Macaulay module.

Lemma 5.2. *Let $\underline{x} = (x_1, \dots, x_d)$ be a system of parameters of M and k a positive integer such that $d - 1 \geq k \geq p_1(M)$. Assume that $\underline{x}$ is a dd-sequence. Then $x_{k+1}, \dots, x_d$ is a standard system of parameters of $M/(x_1, \dots, x_k)M$.*

Proof. Put $M' = M/(x_1, \dots, x_k)M$. Since $\underline{x}$ is a d-sequence, we have

$$\begin{aligned}\chi_1(x_{k+1}, \dots, x_d; M') &= \ell(M'/(x_{k+1}, \dots, x_d)M') - e(x_{k+1}, \dots, x_d; M') \\ &= \ell(M/\underline{x}M) - e(\underline{x}; M) \\ &= \chi_1(\underline{x}; M).\end{aligned}$$

Otherwise, the function $\chi_1(\underline{x}(\underline{n}); M)$ does not depend on $n_{k+1}, \dots, n_d$ by Theorem 1.3. Therefore

$$\begin{aligned}\chi_1(x_{k+1}^{n_{k+1}}, \dots, x_d^{n_d}; M') &= \chi_1(x_1, \dots, x_k, x_{k+1}^{n_{k+1}}, \dots, x_d^{n_d}; M) \\ &= \chi_1(\underline{x}; M) \\ &= \chi_1(x_{k+1}, \dots, x_d; M'),\end{aligned}$$

for all $n_{k+1}, \dots, n_d$. So $x_{k+1}, \dots, x_d$ is a standard system of parameters of M' as required. $\square$

Proof of Theorem 1.4. Since the system of parameters $\underline{x}$ is a dd-sequence, it is not difficult to show by using Lemma 4.2 that

$$\ell(H_{\mathfrak{m}}^0(M/(x_1^{n_1}, \dots, x_i^{n_i})M)) = \sum_{j=0}^i (-1)^{i-j} \binom{d-j-1}{d-i-1} \ell(H_{d-j}(\underline{x}(\underline{n}); M)).$$

Therefore $\ell(H_{\mathfrak{m}}^0(M/(x_1^{n_1}, \dots, x_i^{n_i})M))$ is a polynomial in $\underline{n}$ by Corollary 4.3. For $k \geq p_1(M)$ and positive integers $n_1, \dots, n_k$, we set $M_{k(\underline{n})} = M/(x_1^{n_1}, \dots, x_k^{n_k})M$. Then $M_{k(\underline{n})}$ is a generalized Cohen-Macaulay module and $x_{k+1}, \dots, x_d$ is a standard system of parameters of $M_{k(\underline{n})}$ by Lemma 5.2. Hence, it follows from Lemma 5.1 that

$$\begin{aligned}\ell(H_{\mathfrak{m}}^i(M_{k(\underline{n})})) &= \sum_{j=0}^i (-1)^{i-j} \binom{i}{j} \ell(H_{\mathfrak{m}}^0(M_{k(\underline{n})}/(x_{k+1}, \dots, x_{k+j})M_{k(\underline{n})})) \\ &= \sum_{j=0}^i (-1)^{i-j} \binom{i}{j} \ell(H_{\mathfrak{m}}^0(M/(x_1^{n_1}, \dots, x_k^{n_k}, x_{k+1}, \dots, x_{k+j})M))\end{aligned}$$

is a polynomial in $n_1, \dots, n_k$ for all $i = 0, 1, \dots, d - k - 1$. $\square$

Corollary 5.3. *Let $\underline{x} = (x_1, \dots, x_d)$ be a system of parameters of M . If $\underline{x}$ is a dd-sequence, then $\ell(H_k(x_1^{n_1}, \dots, x_i^{n_i}; M))$ is a polynomial in $n_1, \dots, n_{i-k}$ for all $i = 1, \dots, d$ and $k = 1, \dots, i$.*

Proof. We obtain from the proof of Theorem 1.4 that $\ell(H_{\mathfrak{m}}^0(M/(x_1^{n_1}, \dots, x_i^{n_i})M))$ is a polynomial in $n_1, \dots, n_i$. Therefore $\ell(H_k(x_1^{n_1}, \dots, x_i^{n_i}; M))$ is also a polynomial in $n_1, \dots, n_{i-k}$ by Lemma 4.2. $\square$

6 Sequentially Cohen-Macaulay modules

The concept of sequentially Cohen-Macaulay module was introduced by Stanley [22] for graded modules. Here we define this notion for the local case (see [10], [20]).

Definition 6.1. (i) A filtration $M_0 \subset M_1 \subset \dots \subset M_{t-1} \subset M_t = M$ of submodules of M is called the *dimension filtration* of M if $M_0 = H_{\mathfrak{m}}^0(M)$ and M_{i-1} is the largest submodule of M_i which has dimension strictly less than $\dim M_i$ for all $i = 1, \dots, t$.
(ii) The module M is said to be a *sequentially Cohen-Macaulay module* if each quotient M_i/M_{i-1} is Cohen-Macaulay for all $i = 1, \dots, t$.

Remark 6.2. (i) Because M is Noetherian, the dimension filtration of M always exists and it is unique. Moreover, let $0 = M_0 \subset M_1 \subset \dots \subset M_t = M$ be a dimension filtration of M with $\dim M_i = d_i$. Then we have

$$M_i = \bigcap_{\dim R/\mathfrak{p}_j > d_i} Q_j,$$

for all $i = 0, 1, \dots, t-1$, where $0 = \bigcap_{j=1}^n Q_j$ is a reduced primary decomposition of 0 in M with Q_j is $\mathfrak{p}_j$ -primary for $j = 1, \dots, n$.

(ii) Let $0 = N_0 \subset N_1 \subset \dots \subset N_t = M$ be a filtration of submodules of M . Suppose that each quotient N_i/N_{i-1} is Cohen-Macaulay and $\dim N_1/N_0 < \dim N_2/N_1 < \dots < \dim N_t/N_{t-1}$. Then it was shown in [10] (see also [20]) that this filtration is just the dimension filtration of M and therefore M is sequentially Cohen-Macaulay.

To prove Theorem 1.5 we need some auxiliary lemmata

Lemma 6.3. *Let $M_0 \subset M_1 \subset \dots \subset M_{t-1} \subset M_t = M$ be the dimension filtration of M with $\dim(M_i) = d_i$ and $\underline{x} = (x_1, \dots, x_d)$ a system of parameters of M which is a dd-sequence. Then $M_{t-1} = 0 :_M x_d$.*

Proof. Since $\underline{x}$ is a dd-sequence, $x_d^d \in \mathfrak{a}(M)$ by Corollary 3.9. Therefore $x_d^d \in \text{Ann}_R(0 :_M x)$ for every parameter element x of M by [19, Satz 2.4.5]. By virtue of the Prime Avoidance Theorem we can choose now a parameter element x such that $x \in \text{Ann}_R(M_{t-1})$. Then it follows from the definition of the dimension filtration that

$$M_{t-1} \subseteq 0 :_M x \subseteq 0 :_M x_d^d = 0 :_M x_d \subseteq M_{t-1}.$$

Thus $M_{t-1} = 0 :_M x_d$ as required. $\square$

Lemma 6.4. *Let $M_0 \subset M_1 \subset \cdots \subset M_{t-1} \subset M_t = M$ be the dimension filtration of M with $\dim M_i = d_i$. Assume that M is sequentially Cohen-Macaulay and $x \in \text{Ann}(M_{t-1})$ is a parameter element of M . Then M/xM is a sequentially Cohen-Macaulay module with the dimension filtration*

$$\frac{M_0 + xM}{xM} \subset \frac{M_1 + xM}{xM} \subset \cdots \subset \frac{M_s + xM}{xM} \subset \frac{M}{xM},$$

where $s = t - 2$ if $d_{t-1} = d - 1$ and $s = t - 1$ if $d_{t-1} < d - 1$.

Proof. Since $x \in \text{Ann}(M_{t-1})$, $M_{t-1} \subseteq 0 :_M x^n$ for all $n > 0$ and therefore $M_{t-1} = 0 :_M x^n$ by the definition of the dimension filtration. Then it is easy to check that $M_i \cap xM \subseteq M_{t-1} \cap xM = 0 :_M x \cap xM = 0$ for all $i \leq t - 1$. So we get $\frac{M_i + xM}{xM} \cong M_i$. Thus $\frac{(M_i + xM)/xM}{(M_{i-1} + xM)/xM} \cong M_i/M_{i-1}$ is Cohen-Macaulay for all $i \leq t - 1$. Therefore $(M_s + xM)/xM$ is sequentially Cohen-Macaulay with the dimension filtration

$$\frac{M_0 + xM}{xM} \subset \frac{M_1 + xM}{xM} \subset \cdots \subset \frac{M_s + xM}{xM}$$

by Remark 6.2. Since $\dim(M_s + xM)/xM = d_s < d$, it remains to prove that the module $\frac{M/xM}{(M_s + xM)/xM}$ is a Cohen-Macaulay module. In fact, if $\dim(M_{t-1}) < d - 1$, i. e. $s = t - 1$, we deduce that

$$\frac{M/xM}{(M_{t-1} + xM)/xM} \cong \frac{M/M_{t-1}}{x(M/M_{t-1})}$$

is Cohen-Macaulay since M/M_{t-1} is Cohen-Macaulay. Assume that $\dim(M_{t-1}) = d - 1$, i. e. $s = t - 2$. We have a short exact sequence,

$$0 \longrightarrow \frac{(M_{t-1} + xM)/xM}{(M_{t-2} + xM)/xM} \longrightarrow \frac{M/xM}{(M_{t-2} + xM)/xM} \longrightarrow \frac{M}{M_{t-1} + xM} \longrightarrow 0.$$

where $\frac{(M_{t-1} + xM)/xM}{(M_{t-2} + xM)/xM} \cong M_{t-1}/M_{t-2}$ and $M/(M_{t-1} + xM) \cong \frac{M_t/M_{t-1}}{x(M_t/M_{t-1})}$ are Cohen-Macaulay modules of dimension $d - 1$. So $\frac{M/xM}{(M_{t-2} + xM)/xM}$ is a Cohen-Macaulay module and the lemma is proved. $\square$

Proof of Theorem 1.5

(a) : The implication (i) $\Rightarrow$ (ii) is trivial.

(ii) $\Rightarrow$ (iii) By the assumption $x_1, \dots, x_{d_i}$ is a system of parameters of M_i , so is of $\overline{M}_i = M_i/M_{i-1}$. Since $\overline{M}_i$ is Cohen-Macaulay, we get by [20, Theorem 5.2, a)] that

$$\ell(M/\underline{x}M) = \sum_{i=0}^t \ell(\overline{M}_i/\underline{x}\overline{M}_i) = \sum_{i=0}^t e(x_1, \dots, x_{d_i}; \overline{M}_i) = \sum_{i=0}^t e(x_1, \dots, x_{d_i}; M_i).$$

So

$$\chi_1(\underline{x}; M) = \sum_{i=0}^{t-1} e(x_1, \dots, x_{d_i}; M_i)$$

and (iii) follows since the system of parameters $\underline{x}(\underline{n}) = (x_1^{n_1}, \dots, x_d^{n_d})$ is also satisfied the assumption of (ii)

(iii) $\Rightarrow$ (iv) is straightforward by Theorem 1.2.

(iv) $\Rightarrow$ (i) We proceed by induction on $d = \dim M$. There is nothing to do for the case $d = 1$ since $\underline{x} = (x_1)$ is a strong d-sequence. Assume that $d > 1$. Then $M_{t-1} = 0 :_M x_d$ by Lemma 6.3. Therefore $x_d \in \text{Ann } M_{t-1}$ and $M_{t-1} \cap x_d M = 0$. Put $M' = M/x_d M$. We obtain by Lemma 6.4 that $M' = M/x_d M$ is sequentially Cohen-Macaulay with the dimension filtration

$$\frac{M_0 + x_d M}{x_d M} \subset \frac{M_1 + x_d M}{x_d M} \subset \dots \subset \frac{M_s + x_d M}{x_d M} \subset \frac{M}{x_d M} = M', \quad (*)$$

where $s = t - 2$ if $d_{t-1} = d - 1$ and $s = t - 1$ if $d_{t-1} < d - 1$. Let $d_{t-1} < d - 1$, i. e. $s = t - 1$ in the filtration (*). Since $(x_1, \dots, x_{d-1})$ is a dd-sequence of M' , it follows from the inductive hypotheses for all $i \leq t - 1$ that

$$(M_i + x_d M) \cap (x_{d_i+1}, \dots, x_{d-1}, x_d) M = x_d M.$$

Thus

$$M_i \cap (x_{d_i+1}, \dots, x_d) M \subseteq M_i \cap x_d M \subseteq M_{t-1} \cap x_d M = 0$$

and the implication follows. Since $M_{t-1} \cap x_d M = 0$, the statement is clear for the case $d_{t-1} = d - 1$, i. e. $s = t - 2$ in the filtration (*).

(b): From the short exact sequence $0 \rightarrow M_{t-1} \rightarrow M \rightarrow M/M_{t-1} \rightarrow 0$, we get a long exact sequence of Koszul homology modules

$$\begin{aligned} \dots \rightarrow H_{k+1}(\underline{x}; M/M_{t-1}) \rightarrow H_k(\underline{x}; M_{t-1}) \rightarrow \\ H_k(\underline{x}; M) \rightarrow H_k(\underline{x}; M/M_{t-1}) \rightarrow \dots \end{aligned}$$

Since M/M_{t-1} is Cohen-Macaulay, $H_k(\underline{x}; M/M_{t-1}) = 0$ for all $k > 0$. Therefore $H_k(\underline{x}; M_t) \cong H_k(\underline{x}; M_{t-1})$. So $\chi_k(\underline{x}; M) = \chi_k(\underline{x}; M_{t-1})$. For a positive integer $n \leq d$ we set $\underline{x}_n = (x_1, \dots, x_n)$. Since $x_n M_{t-1} = 0$ for all $n > d_{t-1}$ by the hypothesis, $x_n H_i(\underline{x}_{n-1}; M_{t-1}) = 0$ for all i . Thus, from the long exact sequence

$$\dots \rightarrow H_{i+1}(\underline{x}_n; M_{t-1}) \rightarrow H_i(\underline{x}_{n-1}; M_{t-1}) \xrightarrow{*x_n} H_i(\underline{x}_{n-1}; M_{t-1}) \rightarrow \dots$$

we get the short exact sequences,

$$0 \rightarrow H_{i+1}(\underline{x}_{n-1}; M_{t-1}) \rightarrow H_{i+1}(\underline{x}_n; M_{t-1}) \rightarrow H_i(\underline{x}_{n-1}; M_{t-1}) \rightarrow 0.$$

Therefore we can show by recursive method on $n = d, \dots, d_{t-1} + 1$ that

$$\ell(H_i(\underline{x}; M_{t-1})) = \sum_{j=0}^i \binom{d_t - d_{t-1}}{j} \ell(H_{i-j}(\underline{x}_{d_{t-1}}; M_{t-1}))$$

for all $i \geq 1$. Then for $k > 0$ we have

$$\begin{aligned}
\chi_k(\underline{x}; M) &= \sum_{i=k}^{d_t} (-1)^{i-k} \ell(H_i(\underline{x}; M_{t-1})) \\
&= \sum_{i=k}^{d_t} (-1)^{i-k} \sum_{j=0}^i \binom{d_t - d_{t-1}}{j} \ell(H_{i-j}(\underline{x}_{d_{t-1}}; M_{t-1})) \\
&= \sum_{j=0}^{k-1} \binom{d_t - d_{t-1}}{j} \sum_{i=k}^{d_t} (-1)^{i-k} \ell(H_{i-j}(\underline{x}_{d_{t-1}}; M_{t-1})) \\
&\quad + \sum_{j=k}^{d_t} \binom{d_t - d_{t-1}}{j} \sum_{i=j}^{d_t} (-1)^{i-k} \ell(H_{i-j}(\underline{x}_{d_{t-1}}; M_{t-1})) \\
&= \sum_{j=0}^{k-1} \binom{d_t - d_{t-1}}{j} \chi_{k-j}(\underline{x}_{d_{t-1}}; M_{t-1}) \\
&\quad + (-1)^k \left(\sum_{j=k}^{d_t} (-1)^j \binom{d_t - d_{t-1}}{j} \right) e(\underline{x}_{d_{t-1}}; M_{t-1}).
\end{aligned}$$

Now, we can obtain the formular of the statement (b) by induction on t . Hence the proof of Theorem 1.5 is complete. $\square$

Note that a system of parameters $\underline{x} = (x_1, \dots, x_d)$, which is satisfied the condition (ii) of Theorem 1.5, is just called by P. Schenzel (see [20, Definition 2.5]) a distinguished system of parameters. Since there always exists a distinguished system of parameters on any finitely generated module, it follows immediately from Theorem 1.5, (a) the existence of system of parameters, which is a dd-sequence, on a sequentially Cohen-Macaulay module.

Corollary 6.5. *Any sequentially Cohen-Macaulay module admits a system of parameters which is a dd-sequence.*

An other immediate consequence of Theorem 1.5, (b) is to present a relation between the invariants $p_k(M)$, ($0 \leq k \leq d$) and the dimension $d_i = \dim M_i$ as follows.

Corollary 6.6. *Suppose that M is a sequentially Cohen-Macaulay module. Then for all $i = 0, \dots, d$*

$$p_k(M) = d_i = \dim(R/\mathfrak{a}_{d_i}(M))$$

for $d - d_{i+1} < k \leq d - d_i$, where $\mathfrak{a}_k(M) = \text{Ann}(H_{\mathfrak{m}}^k(M))$.

References

- [1] M. Auslander and D. A. Buchsbaum, Codimension and multiplicity, *Ann. Math.* **68** (1958), 625-657.
- [2] M. P. Brodmann and R. Y. Sharp, *Local cohomology: an algebraic introduction with geometric applications*, Cambridge University Press, 1998.
- [3] W. Bruns and J. Herzog, *Cohen Macaulay rings*, Cambridge University Press, 1993.
- [4] N. T. Cuong, On the length of the powers of systems of parameters in local ring, *Nagoya Math. J.* **120** (1990), 77-88.
- [5] N. T. Cuong, On the dimension of the non-Cohen-Macaulay locus of local rings admitting dualizing complexes, *Math. Proc. Camb. Phil. Soc.* **109** (1991), 479-488.
- [6] N. T. Cuong, On the least degree of polynomials bounding above the differences between lengths and multiplicities of certain systems of parameters in local ring, *Nagoya Math. J.* **125** (1992), 105-114.
- [7] N. T. Cuong, p-standard systems of parameters and p-standard ideals in local rings, *Acta Math. Vietnamica* **20**(1) (1995), 145-161.
- [8] N. T. Cuong and V. T. Khoi, On the partial Euler-Poincaré characteristics of certain systems of parameters in local ring, *Math. Z.* **222** (1996), 383-390.
- [9] N. T. Cuong and N. D. Minh, On the length of Koszul homology and generalized fractions, *Math. Proc. Camb. Phil. Soc.* **120** (1996), 31-42.
- [10] N. T. Cuong and L. T. Nhan, Pseudo Cohen-Macaulay and pseudo generalized Cohen-Macaulay modules, *J. Algebra* **267** (2003), 156-177.
- [11] J-L. Garcia Roig, On polynomial bounds for the of Koszul homology of certain multiplicity systems, *J. London Math. Soc.* **34**(2) (1986), 411-416.
- [12] J-L. Garcia Roig and D. Kirby, On the Koszul homology modules for the powers of multiplicity systems, *Mathematika* **33** (1986), 96-101.
- [13] S. Goto and K. Yamagishi, The theory of unconditioned strong d-sequences and modules of finite local cohomology, preprint (unpublished).
- [14] M. Herrmann, S. Ikeda and U. Orbanz, *Equimultiplicity and Blowing up*, Springer-Verlag, Berlin-Heidelberg-New York 1988.
- [15] C. Huneke, Theory of d-sequences and powers of ideals, *Adv. in Math.* **46** (1982), 249-279.
- [16] T. Kawasaki, On Macaulayfication of Noetherian schemes, *Trans. Amer. Math. Soc.* **352**, no. 6 (2000), 2517-2552.

- [17] T. Kawasaki, On arithmetic Macaulayfication of local rings, *Trans. Amer. Math. Soc.* **354**, no. 1 (2002), 123-149.
- [18] H. Matsumura, *Commutative ring theory*, Cambridge University Press, 1986.
- [19] P. Schenzel, *Dualisierende Komplexe in der lokalen Algebra und Buchsbaum-Ringe*, Lecture Notes in Math. **907**, Springer-Verlag, Berlin-Heidelberg- New York 1982.
- [20] P. Schenzel, *On the dimension filtration and Cohen-Macaulay filtered modules*, Proc. of the Ferrara Meeting in honour of Mario Fiorentini, University of Antwerp, Wilrijk, Belgium, 1998, pp. 245-264.
- [21] J. P. Serre, *Algèbre locale. Multiplicités*, 3rd ed., Lecture Notes in Math. **11**, Springer-Verlag, Berlin-Heidelberg-New York 1975.
- [22] R. P. Stanley, *Combinatorics and Commutative Algebra*, Second edition, Birkhäuser Boston 1996.
- [23] J. Stückrad and W. Vogel, *Buchsbaum rings and applications*, Springer-Verlag, Berlin-Heidelberg-New York 1986.
- [24] N. V. Trung, Toward a theory of generalized Cohen-Macaulay modules, *Nagoya Math. J.* **102** (1986), 1-49.